

CORRECTION

Open Access

Correction: Alcohol-related brief intervention in patients treated for opiate or cocaine dependence: a randomized controlled study

Nelson Feldman*, Anne Chatton, Riaz Khan, Yasser Khazaal and Daniele Zullino

Correction

After publication of this article, it came to our attention that the published manuscript contained errors in the table three (Table 1), Table four (Table 2), Table five (Table 3) and Table six (Table 4), Table seven (Table 5). The original published article contains old tables from the first version of the manuscript sent for initial consideration for publication.

This correction article contains the tables revised after peer review of the manuscript.

The authors and Publisher apologize to the readers for the inconvenience caused.

This correction article contains the revised tables from the original article [1] Tables one and two remain unchanged.

Table 1 AUDIT scores and number of weekly alcohol drinks by group

	CONTROL GROUP			INTERVENTION GROUP			CONTROL AND INTERVENTION GROUPS		
	T0	T3	T9	T0	T3	T9	T0	T3	T9
AUDIT, Mean (±SD)	16.6 (8.2)	12.4 (7.5)	11.6 (7.3)	16.9 (7.7)	13 (6.2)	13.9 (8.1)	16.8 (7.9)	12.7 (6.8)	12.8 (7.8)
Number of drinks/week, Mean (±SD)	23.8 (9.5)	16.3 (12.6)	18.7 (13.2)	25.3 (14.3)	17 (11.6)	18.4 (10.4)	24.6 (12.3)	16.7 (12)	18.5 (11.7)

Table 2 AUDIT scores and number of alcohol drinks per week in excessive drinkers and alcohol dependents

	EXCESSIVE DRINKERS			ALCOHOL DEPENDENTS		
	T0	T3	T9	T0	T3	T9
AUDIT, Mean (± SD)	9.4 (1.2)	10.6 (6)	9.4 (5.8)	21.9 (6.2)	14.2 (7)	15.2 (8.1)
Number of drinks per week, Mean (± SD)	19.4 (9.2)	13.4 (7)	15.1 (8.3)	28.2 (12.9)	19 (14.1)	20.9 (13.1)

Table 3 AUDIT scores according to group and gender

	CONTROL GROUP			INTERVENTION GROUP			CONTROL AND INTERVENTION GROUPS		
	T0	T3	T9	T0	T3	T9	T0	T3	T9
Men	17.6 (9)	13.4 (7.4)	13.4 (6.8)	17.3 (7.9)	13.5 (6.3)	14.3 (8.7)	17.4 (8.4)	13.4 (6.8)	13.6 (8)
Women	14.2 (5.4)	10 (7.4)	9 (7.3)	16 (6.9)	11.3 (5.7)	12.6 (6)	15.1 (6.2)	10.7 (6.5)	10.8 (6.8)

Table 4 Number of alcohol drinks per week according to group and gender

	CONTROL GROUP			INTERVENTION GROUP			CONTROL AND INTERVENTION GROUPS		
	T0	T3	T9	T0	T3	T9	T0	T3	T9
Men	23.9 (10.3)	17.7 (13.6)	19.8 (14.4)	26.7 (15.3)	17.7 (12.9)	18.5 (11)	25.5 (13.3)	17.7 (13.1)	19.1 (12.5)
Women	23.4 (7.7)	13.1 (9.3)	16 (9.9)	21.2 (10.1)	14.9 (6.5)	18 (8.8)	22.3 (8.9)	14 (7.9)	17 (9.3)

* Correspondence: Nelson.feldman@hcuge.ch
 Geneva University Hospitals, Geneva, Switzerland

Table 5 Evolution of frequency of alcohol consumption by treatment group

	CONTROL GROUP			INTERVENTION GROUP		
	T0-T3	T3-T9	T0-T9	T0-T3	T3-T9	T0-T9
Increased or unchanged frequency of alcohol use: % failure (n)	68.6 (35)	78.4 (40)	70.6 (36)	64.4 (38)	71.2 (42)	67.8 (40)
Decreased frequency of alcohol use: % success (n)	31.4 (16)	21.6 (11)	29.4 (15)	35.6 (21)	28.8 (17)	32.2 (19)

Received: 31 July 2013 Accepted: 31 July 2013

Published: 1 August 2013

Reference

1. Feldman N, Chatton A, Khan R, Khazaal Y, Zullino D: Alcohol-related brief intervention in patients treated for opiate or cocaine dependence: a randomized controlled study. *Substance Abuse Treatment, Prevention, and Policy* 2011, **6**:22.

doi:10.1186/1747-597X-8-28

Cite this article as: Feldman et al.: Correction: Alcohol-related brief intervention in patients treated for opiate or cocaine dependence: a randomized controlled study. *Substance Abuse Treatment, Prevention, and Policy* 2013 **8**:28.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

